


S1: (arrow marks) PRp = 95% +/-2% @ 1064 nm + PRa = 95% +/-2% @ 532 nm, AOI 45 deg  
S2: ARp < 0.2% @ 1064 nm + ARa < 0.8% @ 532 nm, AOI




PO8346 Ra i45

Fig. 1.

SIDE MEASURED: S1 only (grinded witness sample)

COMMENTS:

S1: (arrow marks) PRp = 95% +/-2% @ 1064 nm + PRa = 95% +/-2% @ 532 nm, AOI 45 deg  
S2: ARp < 0.2% @ 1064 nm + ARa < 0.8% @ 532 nm, AOI




PO8346 Ra i45

Fig. 2.

SIDE MEASURED: S2 only (grinded witness sample)

COMMENTS:

S1: (arrow marks) PRp = 95% +/-2% @ 1064 nm + PRa = 95% +/-2% @ 532 nm, AOI 45 deg  
S2: ARp < 0.2% @ 1064 nm + ARa < 0.8% @ 532 nm, AOI




PO8346 Rp i45

Fig. 3.

SIDE MEASURED: S1 only (grinded witness sample)

COMMENTS:

S1: (arrow marks) PRp = 95% +/-2% @ 1064 nm + PRa = 95% +/-2% @ 532 nm, AOI 45 deg  
S2: ARp < 0.2% @ 1064 nm + ARa < 0.8% @ 532 nm, AOI



PO8346 Rp i45

Fig. 4.

SIDE MEASURED: S2 only (grinded witness sample)

COMMENTS: